

D0 and Tevatron results on the standard model Higgs boson using the full Run 2 data

*Lidija Živković,
LPNHE, Paris VI & VII
on behalf of the*



and




collaborations

*Joint Experimental-Theoretical Seminar,
Fermilab
22 March 2013*


Outline

- Current status
- Overview of the D0 search channels
- D0 results
 - Combinations of different channels
- Tevatron results
 - D0 and CDF combinations
 - Constraints on couplings




Introduction

- SM Higgs boson mass is constrained indirectly through precision measurements
 - self-energy corrections to the W mass depend on the mass of the top quark and Higgs boson, *which are both precisely measured at Tevatron*







- Global SM electroweak fits provide upper limit
 - The best fit gives $m_H = 94^{+29}_{-24}$ GeV
 - Limit from fit $m_H < 152$ GeV



Historical perspective



- LEP (1989 - 2000): $m_H > 114.4 \text{ GeV}@95\% \text{ CL}$
- At hadron colliders:
- Tevatron Run II (2002-2011):
 - First post-LEP exclusion (2009)
 - First evidence of a Higgs-like particle decaying to a pair of b-quarks (July 2012)
- LHC (2009 - 2012):
 - Excluded wide mass range (111 - 122 GeV and 127 - 600 GeV)
 - Discovered the new Higgs-like boson mainly through $\gamma\gamma$ and ZZ decays (July 2012)



Current situation

- LHC (2009 - 2012):
 - Since July 2012 progress in each channel
 - Observation confirmed in bosonic channel
 - ATLAS: $m_H = 125.5 \pm 0.2 \text{ (stat)} -0.6+0.5 \text{ (sys)} \text{ GeV}$
 - CMS: $m_H = 125.8 \pm 0.4 \text{ (stat)} \pm 0.4 \text{ (sys)} \text{ GeV}$
 - $H \rightarrow bb$ with $\sim 18 \text{ fb}^{-1}$ data deficit at Atlas and $\sim 2.2 \text{ s.d. excess}$ at CMS

*As presented at
Moriond and Aspen*




Tevatron Data Taking



Run II Integrated Luminosity

19 April 2002 - 30 September 2011




Tevatron Data Taking





Run II Integrated Luminosity

19 April 2002 - 30 September 2011





Production at Tevatron ...

- Dominant production is **gluon-gluon fusion (ggH)**
- Significant contribution from **associated production (VH)**



... and Decay


- Dominant decay to:
 - bb for $m_H < 135 \text{ GeV}$
 - WW for $m_H > 135 \text{ GeV}$




Backgrounds

- We model background processes with Alpgen+Pythia, Pythia and CompHEP
- Normalized with the highest order cross section available (NLO or better)


Z+jets
 $\sigma \sim O(10^3 \text{ pb})$



Multijet (QCD)
from data




W+jets
 $\sigma \sim O(10^3 \text{ pb})$




Higgs signal
 $\sigma \sim O(0.1 \text{ pb})$




tt
 $\sigma \sim O(8 \text{ pb})$




diBoson
 $\sigma \sim O(10 \text{ pb})$




Single top
 $\sigma \sim O(3 \text{ pb})$




How did we search?



How did we search?



How did we search?



- Extract tiny signal from huge background
- Use efficient and well understood triggers
- Optimize lepton ID, use multivariate techniques (MVA) to identify leptons and various lepton categories
- Optimize b-id, use MVA and multiple b-tagging categories
- Use advanced MVA techniques to further separate signal from background
- Validate search with measurement of known SM processes

Overview of the D0 results

CDF results presented by T. Junk on January 18th:

http://theory.fnal.gov/jetp/talks/wc_trj_cdfhiggs_18jan_pub.pdf

Overview of the channels at D0

DØ	Luminosity (fb ⁻¹)	M_H (GeV)	Reference
$WH \rightarrow \ell \nu b \bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121804 (2012); Acc by PRD arXiv:1301.6122
$ZH \rightarrow \ell \ell b \bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121803 (2012); Sub to PRD arXiv:1303.3276
$ZH \rightarrow \nu \bar{\nu} b \bar{b}$	9.5	100–150	Phys. Lett. B 716, 285 (2012)
$H \rightarrow W^+ W^- \rightarrow \ell^+ \nu \ell^- \bar{\nu}$	9.7	100–200	Acc by PRD arXiv:1301.1243
$H + X \rightarrow WW \rightarrow \mu^\pm \tau_h^\mp + \leq 1\text{jet}$	7.3	155–200	Phys. Lett. B 714, 237 (2012)
$H \rightarrow W^+ W^- \rightarrow \ell \nu q' \bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow e e \mu / \mu \mu e + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow e^\pm \mu^\pm + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow \ell \nu q' \bar{q} q' \bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow \tau_h \tau_h \mu + X$	8.6	100–150	Sub to PRD arXiv:1302.5723
$H + X \rightarrow \ell \tau_h j j$	9.7	105–150	Acc by PRD arXiv:1211.6993
$H \rightarrow \gamma \gamma$	9.7	100–150	Acc by PRD, arXiv:1301.5358




- D0 combination:
 $H \rightarrow b \bar{b}$:
 - Phys. Rev. Lett. 109, 121802 (2012)
 All channels:
 - Submitted to PRD arXiv:1303.0823
- All new papers will be in a single issue of PRD

$$H \rightarrow \gamma\gamma$$

$$H \rightarrow \gamma\gamma$$

New since
ICHEP'12

- Updated data quality requirement
- Narrow resonance on top of a smoothly falling background in the $m_{\gamma\gamma}$ spectrum: Measured mass resolution 3.1 GeV @ $m_H=125$ GeV
- Multiple stages of MVA:
 - Neural Network to select loose photons, and then to define two independent samples, γ -enriched and jet-enriched
 - Boosted Decision Trees to further separate signal from backgrounds
- Exp. (obs.) sensitivity @125 GeV: 8.7*SM (12.8*SM)






$$H \rightarrow \tau\tau$$

$$H \rightarrow \tau\tau$$

New since
ICHEP'12


- Combine analyses with $H \rightarrow \tau\tau$:
 - $VH \rightarrow \mu\tau\tau$ - added ~20% more data, optimized
 - $H+X \rightarrow l\tau + 2 \text{ jets}$ - added
- They include some contribution from $H \rightarrow WW$
 - Use dedicated MVA to separate different contributions
- Exp. (obs.) @125 GeV: 7.25 (10.84)*SM
(~60% improvement from ICHEP'12)



$$H \rightarrow WW$$

$$H \rightarrow WW$$


- The most sensitive channel for $m_H > 135 \text{ GeV}$
- Split according to the decay mode of the W
 - Dilepton channels have low branching ratios thus low yield, but also low backgrounds
 - Semileptonic channels must contend with large V +jets backgrounds
- Split according to the production mode - ggH , VBF and VH
 - Split opposite-sign dilepton and semileptonic channels into different jet multiplicities
 - Include a search for the same-sign leptons, where one originates from associated W
 - Include final states with three leptons




Associated production with W , $WH \rightarrow WWW$ is important for coupling measurement; It probes coupling to the W boson only!

New since
ICHEP'12



$$H \rightarrow WW \rightarrow l\nu l\nu$$



- Refined selection; splitting to WW enriched and depleted regions for $H \rightarrow WW \rightarrow e\nu\mu\nu$
- Signatures:
 - Two isolated high p_T leptons, large E_T , small $\Delta\phi(l\bar{l})$, small $\Delta R(l\bar{l})$
- \Rightarrow Lepton identification improved over years to allow for more efficient selection
- Use multiple MVA to reject different backgrounds






- Remove most of the Z(+jets) \rightarrow ll
- Use dedicated MVA to separate samples into WW enriched and WW depleted regions



Validation with diboson measurement





- Use identical selection and similar MVA
 - Use WW process as a signal in training
- Measured cross section: $(1.02 \pm 0.06) * SM$



Other $H \rightarrow WW$ searches

New since
ICHEP'12

- Included semileptonic decays of W , and also associated production:
 - $H \rightarrow WW \rightarrow l\nu jj$ - added ~80% of data
 - $VH \rightarrow VWW \rightarrow l\nu + 4 \text{ jets}$ - optimized
 - $VH \rightarrow VVV \rightarrow lll + X$ - added ~12% of data
 - $VH \rightarrow VWW \rightarrow e^+\mu^+ + X$ - optimized





$H \rightarrow WW$ result

- Dilepton channel only:
- Excludes (expect):
159-176 (156-172) GeV
- Sensitivity @125 GeV:
exp - 3.4; obs - 4.1

Gain 15%


- Full D0 $H \rightarrow WW$:
- Excludes (expect):
157-178 (155-175) GeV
- Sensitivity @125 GeV
exp - 2.9; obs - 4.6



Log Likelihood Ratio (LLR)

Background-like
outcomes

Signal-like
outcomes




$$LLR = -2 \ln \frac{P(s+b)}{P(b)}$$

P - Poisson likelihood of B or S+B hypothesis

- The separation between LLR_b (background-only hypothesis) and LLR_{s+b} (signal-plus-background hypothesis) provides a measure of the discriminating power of the search
- The width of the LLR_b distribution (1 s.d. and 2 s.d. bands) provides an estimate of how sensitive the analysis is to a signal-like background fluctuation in the data, taking account of the presence of systematic uncertainties
- The value of LLR_{obs} relative to LLR_{s+b} and LLR_b indicates whether the data distribution appears to be more like signal-plus-background or background-only.



$$VH \rightarrow Vbb$$

$VH \rightarrow Vbb$



- $ZH \rightarrow llbb$ - 2 leptons + 2 b-jets
- Modeling of the Z+jets background; rejection of the tt background



- $WH \rightarrow lvbb$ - 1 lepton + MET + 2 b-jets
- Modeling of the W+jets backgrounds
- Modeling and rejection of the multijet backgrounds





- $ZH \rightarrow \nu\nu bb$ - MET + 2 b-jets (contribution from WH also)
- Background modeling and rejection

VH \rightarrow Vbb

- Key ingredients:
 - Lepton, jet and E_T reconstruction
 - Jet energy resolution $\Rightarrow \Delta m/m \sim 15\%$
 - **b-tagging**
 - Multivariate techniques to reject backgrounds





	Before b-tagging	2 loose tags
s/b	1/7000	1/1400





VH \rightarrow Vbb

- Key ingredients:
 - Lepton, jet and E_T reconstruction
 - Jet energy resolution $\Rightarrow \Delta m/m \sim 15\%$
 - **b-tagging**
 - Multivariate techniques to reject backgrounds





	Before b-tagging	2 med tags
s/b	1/7000	1/400





VH \rightarrow Vbb

- Key ingredients:
 - Lepton, jet and E_T reconstruction
 - Jet energy resolution $\Rightarrow \Delta m/m \sim 15\%$
 - **b-tagging**
 - Multivariate techniques to reject backgrounds







	Before b-tagging	2 tight tags
s/b	1/7000	1/200




VH \rightarrow Vbb

- Key ingredients:
 - Lepton, jet and \cancel{E}_T reconstruction
 - Jet energy resolution (15%)
 - b-tagging: eff 50-80%; mis id 1-10%
 - Multivariate techniques to reject backgrounds
 - Either to split into two regions or to remove background



Validation of results






- Measure cross section of the known process with the same final state
 - Smaller cross section for Higgs production (~ 7 times)
 - Diboson signal peaks at lower masses

- Apply similar analysis







- Measured cross section:
 $(0.73 \pm 0.32) \times SM$

MH = 125 GeV	VH \rightarrow Vbb [fb]	VZ \rightarrow Vbb [fb]
vvbb	9	73
lvbb	16	105
llbb	3	24
Total	28	202




VH \rightarrow Vbb results

- Exp (obs) @125 GeV:
- 3.9(4.3)*SM
- Exp (obs) @125 GeV:
- 4.7(4.8)*SM
- Exp (obs) @125 GeV:
- 5.1(7.1)*SM





VH \rightarrow Vbb results

- Expected sensitivity @125 GeV: $2.3 \times \text{SM}$; observed $3.5 \times \text{SM}$



D0 combination

Combining all D0 channels: What does combined signal look like?




- Distribution of $\log_{10}(s/b)$, for the data from all contributing Higgs boson search channels

Sensitivity of the search

Background-like
outcomes


Signal-like
outcomes



- Observe a broad excess between $\sim 115 \text{ GeV}$ and $\sim 145 \text{ GeV}$ consistent with a SM Higgs expectation


p-value for background hypothesis

- p-value for background hypothesis provides information about the consistency with the observed data
- Local p-value distribution for background only expectation:
 - DØ: 1.7 s.d. (@125 GeV)





Signal Strength

- Best fit for the signal, signal strength, is consistent with SM within 1 s.d.
- @125 GeV: $1.40^{+0.92}_{-0.88}$



Signal Strength

- Best fit for the signal, signal strength, is consistent with SM within 1 s.d.
- @125 GeV: $1.40^{+0.92}_{-0.88}$



Combined	$1.40^{+0.92}_{-0.88}$
$H \rightarrow \gamma\gamma$	$4.20^{+4.60}_{-4.20}$
$H \rightarrow W^+W^-$	$1.90^{+1.63}_{-1.52}$
$H \rightarrow \tau^+\tau^-$	$3.96^{+4.11}_{-3.38}$
$H \rightarrow b\bar{b}$	$1.23^{+1.24}_{-1.17}$

Tevatron combination

Overview of the searches

DØ	Luminosity (fb ⁻¹)	M_H (GeV)	Reference
$WH \rightarrow \ell\nu b\bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121804 (2012); Acc by PRD arXiv:1301.6122
$ZH \rightarrow \ell\ell b\bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121803 (2012); Sub to PRD arXiv:1303.3276
$ZH \rightarrow \nu\bar{\nu} b\bar{b}$	9.5	100–150	Phys. Lett. B 716, 285 (2012)
$H \rightarrow W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$	9.7	100–200	Acc by PRD arXiv:1301.1243
$H + X \rightarrow WW \rightarrow \mu^\pm\tau_h^\mp + \leq 1\text{jet}$	7.3	155–200	Phys. Lett. B 714, 237 (2012)
$H \rightarrow W^+W^- \rightarrow \ell\nu q'\bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow e^\pm\mu^\pm + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow \ell\nu q'\bar{q}q'\bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow \tau_h\tau_h\mu + X$	8.6	100–150	Sub to PRD arXiv:1302.5723
$H + X \rightarrow \ell\tau_h jj$	9.7	105–150	Acc by PRD arXiv:1211.6993
$H \rightarrow \gamma\gamma$	9.7	100–150	Acc by PRD, arXiv:1301.5358
CDF			
$WH \rightarrow \ell\nu b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111804 (2012)
$ZH \rightarrow \ell\ell b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111803 (2012)
$ZH \rightarrow \nu\bar{\nu} b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111805 (2012); Acc. by PRD arXiv: 1301.4440
$H \rightarrow W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow WW \rightarrow e\tau_h\mu\tau_h$	9.7	130–200	FERMILAB-PUB-13-029-E, For submission to PRD
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow \tau\tau$	6.0	100–150	Phys. Rev. Lett. 108, 181804 (2012)
$H \rightarrow \gamma\gamma$	10.0	100–150	Phys. Lett. B 717, 173 (2012)
$H \rightarrow ZZ \rightarrow llll$	9.7	120–200	Phys. Rev. D 86 (2012) 072012
$t\bar{t}H \rightarrow WWb\bar{b}b\bar{b}$	9.45	100–150	Phys. Rev. Lett. 109 (2012) 181802
$VH \rightarrow jjb\bar{b}$	9.45	100–150	JHEP 1302 (2013) 004

- CDF combination:

$H \rightarrow b\bar{b}$:

- Phys. Rev. Lett. 109, 111802 (2012)

All channels:

- Submitted to PRD arXiv:1301.6668

- DØ combination:

$H \rightarrow b\bar{b}$:

- Phys. Rev. Lett. 109, 121802 (2012)

All channels:

- Submitted to PRD arXiv:1303.0823

Overview of the searches

DØ	Luminosity (fb ⁻¹)	M_H (GeV)	Reference
$WH \rightarrow \ell\nu b\bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121804 (2012); Acc by PRD arXiv:1301.6122
$ZH \rightarrow \ell\ell b\bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121803 (2012); Sub to PRD arXiv:1303.3276
$ZH \rightarrow \nu\bar{\nu} b\bar{b}$	9.5	100–150	Phys. Lett. B 716, 285 (2012)
$H \rightarrow W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$	9.7	100–200	Acc by PRD arXiv:1301.1243
$H + X \rightarrow WW \rightarrow \mu^\pm\tau_h^\mp + \leq 1\text{jet}$	7.3	155–200	Phys. Lett. B 714, 237 (2012)
$H \rightarrow W^+W^- \rightarrow \ell\nu q'\bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow e^\pm\mu^\pm + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow \ell\nu q'\bar{q}q'\bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow \tau_h\tau_h\mu + X$	8.6	100–150	Sub to PRD arXiv:1302.5723
$H + X \rightarrow \ell\tau$	9.7	100–150	Acc by PRD arXiv:1301.6002
$H \rightarrow \gamma\gamma$			
CDF			
$WH \rightarrow \ell\nu b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111804 (2012)
$ZH \rightarrow \ell\ell b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111803 (2012)
$ZH \rightarrow \nu\bar{\nu} b\bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111805 (2012); Acc. by PRD arXiv: 1301.4440
$H \rightarrow W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow WW \rightarrow e\tau_h\mu\tau_h$	9.7	130–200	FERMILAB-PUB-13-029-E, For submission to PRD
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow \tau\tau$	6.0	100–150	Phys. Rev. Lett. 108, 181804 (2012)
$H \rightarrow \gamma\gamma$	10.0	100–150	Phys. Lett. B 717, 173 (2012)
$H \rightarrow ZZ \rightarrow llll$	9.7	120–200	Phys. Rev. D 86 (2012) 072012
$t\bar{t}H \rightarrow WWb\bar{b}b\bar{b}$	9.45	100–150	Phys. Rev. Lett. 109 (2012) 181802
$VH \rightarrow jjb\bar{b}$	9.45	100–150	JHEP 1302 (2013) 004

Tevatron combination is about to be submitted!

• CDF combination:

$H \rightarrow b\bar{b}$:

- Phys. Rev. Lett. 109, 111802 (2012)

All channels:

- Submitted to PRD arXiv:1301.6668

• DØ combination:

$H \rightarrow b\bar{b}$:

- Phys. Rev. Lett. 109, 121802 (2012)


All channels:

- Submitted to PRD arXiv:1303.0823



Validation with diboson measurement

Validation of results

- Measure cross section of the known process with the same final state
 - Smaller cross section for Higgs production (~ 7 times)
 - Diboson signal peaks at lower masses
- Apply similar analysis
- Measured cross section: $(0.68 \pm 0.21) \times \text{SM}$





MH = 125 GeV	VH \rightarrow Vbb [fb]	VZ \rightarrow Vbb [fb]
vvbb	9	73
lvbb	16	105
llbb	3	24
Total	28	202



Combined results from Tevatron


Decay mode combinations



- $H \rightarrow \gamma\gamma$
 - Expected sensitivity @125 GeV of $\sim 5.9 \cdot \text{SM}$
 - ~ 2 s.d. excess in $H \rightarrow \gamma\gamma$



- $H \rightarrow \tau\tau$
 - Expected sensitivity @125 GeV of $\sim 5.7 \cdot \text{SM}$

Decay mode combinations





- $H \rightarrow WW$:
 - Expected sensitivity @125 GeV of $2.04 \times \text{SM}$
 - Very broad excess consistent with expectations (i.e., lack of mass peak due to escaping neutrinos)

Decay mode combinations




- $VH \rightarrow Vbb$:
 - Expected sensitivity at $m_H \sim 125 \text{ GeV}$ of $1.42 \times \text{SM}$.
 - Broad excess consistent with dijet mass resolution
 - Best fit $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow bb) = 0.19^{+0.08}_{-0.09} \text{ pb @125 GeV}$
 - To be compared with SM: $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow bb) = 0.12 \pm 0.01 \text{ pb}$

Combining all CDF and D0 channels: What does combined signal look like?




- Distribution of $\log_{10}(s/b)$, for the data from all combining Higgs boson decay channels

Result of the SM combination




- Tevatron *excludes* (expect):
90-109 (90-120) GeV and 149-182 (140-184) GeV @95% C.L.
- Exp. (obs) sensitivity @125 GeV: 1.06 (2.44)*SM

Sensitivity of the search



- Observed broad excess in data


Sensitivity of the search



- Observed broad excess in data
 - Consistent with the assumption of the presence of the Higgs boson with a $m_H = 125 \text{ GeV}$ and a cross section of $\sim 1.5 \times \text{SM}$


p-value for background hypothesis

- p-value for background hypothesis provides information about the consistency with the observed data
- Local p-value distribution for background only expectation:
 - 3.1 s.d. (@125 GeV)



Signal Strength

- Best fit for the signal, signal strength, is consistent with SM within 1 s.d.
- @125 GeV: $1.44^{+0.59}_{-0.56}$



Signal strengths for various decays



DØ	Luminosity (fb ⁻¹)	M_H (GeV)	Reference
$WH \rightarrow \ell \nu b \bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121804 (2012); Acc by PRD arXiv:1301.6122
$ZH \rightarrow \ell \ell b \bar{b}$	9.7	90–150	Phys. Rev. Lett. 109, 121803 (2012); Sub to PRD arXiv:1303.3276
$ZH \rightarrow \nu \bar{\nu} b \bar{b}$	9.5	100–150	Phys. Lett. B 716, 285 (2012)
$H \rightarrow W^+ W^- \rightarrow \ell^+ \nu \ell^- \bar{\nu}$	9.7	100–200	Acc by PRD arXiv:1301.1243
$H + X \rightarrow WW \rightarrow \mu^\pm \tau_h^\mp + \leq 1\text{jet}$	7.3	155–200	Phys. Lett. B 714, 237 (2012)
$H \rightarrow W^+ W^- \rightarrow \ell \nu q' \bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow e^\pm \mu^\pm + X$	9.7	100–200	Sub to PRD arXiv:1302.5723
$VH \rightarrow \ell \nu q' \bar{q} q' \bar{q}$	9.7	100–200	Acc by PRD arXiv:1301.6122
$VH \rightarrow \tau_h \tau_h \mu + X$	8.6	100–150	Sub to PRD arXiv:1302.5723
$H + X \rightarrow \ell \tau_h jj$	9.7	105–150	Acc by PRD arXiv:1211.6993
$H \rightarrow \gamma\gamma$	9.7	100–150	Acc by PRD, arXiv:1301.5358
CDF			
$WH \rightarrow \ell \nu b \bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111804 (2012)
$ZH \rightarrow \ell \ell b \bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111803 (2012)
$ZH \rightarrow \nu \bar{\nu} b \bar{b}$	9.45	90–150	Phys. Rev. Lett. 109, 111805 (2012); Acc. by PRD arXiv: 1301.4440
$H \rightarrow W^+ W^- \rightarrow \ell^+ \nu \ell^- \bar{\nu}$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow WW \rightarrow e \tau_h \mu \tau_h$	9.7	130–200	FERMILAB-PUB-13-029-E, For submission to PRD
$VH \rightarrow ee\mu/\mu\mu e + X$	9.7	110–200	FERMILAB-PUB-13-029-E, For submission to PRD
$H \rightarrow \tau\tau$	6.0	100–150	Phys. Rev. Lett. 108, 181804 (2012)
$H \rightarrow \gamma\gamma$	10.0	100–150	Phys. Lett. B 717, 173 (2012)
$H \rightarrow ZZ \rightarrow ll ll$	9.7	120–200	Phys. Rev. D 86 (2012) 072012
$t\bar{t}H \rightarrow WW b\bar{b} b\bar{b}$	9.45	100–150	Phys. Rev. Lett. 109 (2012) 181802
$VH \rightarrow jj b\bar{b}$	9.45	100–150	JHEP 1302 (2013) 004

- $VH \rightarrow Vbb$; $H \rightarrow WW$; $H \rightarrow \tau\tau$; $H \rightarrow \gamma\gamma$;

Signal strength for various decays


- Posterior probability densities for $R = (\sigma \times \mathcal{B})/SM$ from the combinations of all search channels

m_H (GeV/c ²)	125
$R_{\text{fit}}(SM)$	$1.44^{+0.59}_{-0.56}$
$R_{\text{fit}}(H \rightarrow W^+W^-)$	$0.94^{+0.85}_{-0.83}$
$R_{\text{fit}}(H \rightarrow b\bar{b})$	$1.59^{+0.69}_{-0.72}$
$R_{\text{fit}}(H \rightarrow \gamma\gamma)$	$5.97^{+3.39}_{-3.12}$
$R_{\text{fit}}(H \rightarrow \tau^+\tau^-)$	$1.68^{+2.28}_{-1.68}$




Little history of Tevatron results


Data of 2007; up to 2.4 fb^{-1}




Data of 2008; up to 4.2 fb^{-1}




Data of mid 2009; up to 5.4 fb^{-1}




Time



Data of mid 2010; up to 5.9 fb^{-1}



Data of mid 2011; up to 8.6 fb^{-1}






Full data set; up to 10 fb^{-1}

Couplings

Higgs boson couplings to bosons and fermions

- Several production and decay mechanisms contribute to signal rates per channel => interpretation is difficult
- Simplified model, SM-like with the following:
 - Hff couplings are scaled together by κ_f
 - HWW coupling is scaled by κ_W
 - HZZ coupling is scaled by κ_Z
- For some studies, we scale the HWW and HZZ couplings by $\kappa_W = \kappa_Z = \kappa_V$
- Standard Model is recovered if $\kappa_f = \kappa_W = \kappa_Z = 1$



Higgs boson couplings to bosons and fermions

- Follow the prescription from LHC Higgs cross section working group: [arXiv:1209.0040](https://arxiv.org/abs/1209.0040)
- Basic assumptions:
 - There is only one underlying state at $m_H \sim 125 \text{ GeV}$
 - It has negligible width
 - It is a CP even scalar (only allow for modification of coupling strengths, leaving the Lorentz structure of the interaction untouched)
 - No additional invisible or undetected Higgs decay modes

Constraining couplings

- Scale cross sections for each process according to couplings

$$\sigma(gg \rightarrow H) = \sigma_{SM}(gg \rightarrow H)(0.95\kappa_f^2 + 0.05\kappa_f\kappa_V)$$

$$\sigma(VH, VBF) = \sigma_{SM}(VH, VBF)\kappa_V^2$$

- Recompute all Higgs boson decay branching ratios from scaled partial widths

$$\Gamma(H \rightarrow VV) = \Gamma(H \rightarrow VV)_{SM}\kappa_V^2; (V = W, Z)$$

$$\Gamma(H \rightarrow ff) = \Gamma(H \rightarrow ff)_{SM}\kappa_f^2$$

$$\Gamma(H \rightarrow gg) = \Gamma(H \rightarrow gg)_{SM}(0.95\kappa_f^2 + 0.05\kappa_f\kappa_V)$$

$$\Gamma(H \rightarrow \gamma\gamma) = \Gamma(H \rightarrow \gamma\gamma)_{SM}|\alpha\kappa_V + \beta\kappa_f|^2$$

$$BR(H \rightarrow XX) = \frac{\Gamma(H \rightarrow XX)}{\Gamma_{TOT}}$$


$$\alpha=1.28; \beta=-0.21;$$

from Spira et al. arXiv:hep-ph/9504378

=> $H \rightarrow \gamma\gamma$ from destructive interference between the two contributions

- If any of the couplings is negative, interference becomes constructive

=> Larger rate of the $H \rightarrow \gamma\gamma$



Couplings

- Posterior probability distributions

(a) vary κ_W ($\kappa_Z = \kappa_f = 1$)




- A negative sign of κ_W is preferred by the Tevatron data due to the excess in $H \rightarrow \gamma\gamma$
- Best fit: $\kappa_W = -1.27$

(b) vary κ_Z ($\kappa_W = \kappa_f = 1$)

- Searches at the Tevatron are sensitive almost exclusively to $(\kappa_Z)^2$ so the posterior density is nearly symmetric
- Best fit: $\kappa_Z = \pm 1.05$


(c) vary κ_f ($\kappa_W = \kappa_Z = 1$)

- Asymmetry due to $H \rightarrow \gamma\gamma$
- Best fit: $\kappa_f = -2.64$ (large due to the excesses in $H \rightarrow \gamma\gamma$ and $VH \rightarrow Vbb$)




Couplings

- Both κ_W and κ_Z vary independently
 - κ_f integrated over
 - Best fit: $(\kappa_W, \kappa_Z) = (1.25, \pm 0.90)$
- The point $(\kappa_W, \kappa_Z) = (0, 0)$ corresponds to no Higgs boson production or decay in the most sensitive search modes at the Tevatron and is not included within the 95% C.L. region due to the significant excess of events in the SM Higgs boson searches @ 125 GeV




Couplings

- Probe $SU(2)_V$ custodial symmetry by measuring the ratio $\lambda_{WZ} = \kappa_W / \kappa_Z$
 - Measure $\theta_{WZ} = \tan^{-1}(\kappa_Z / \kappa_W) = \tan^{-1}(1/\lambda_{WZ})$
 - Measure: $|\theta_{WZ}| = 0.68^{+0.21}_{-0.41} \rightarrow \lambda_{WZ} = 1.24^{+2.34}_{-0.42}$



Couplings

- Assuming that custodial symmetry holds, $\lambda_{WZ} = 1$, allow both κ_V and κ_f to vary
- Asymmetry is from the excesses in the $H \rightarrow \gamma\gamma$
- Two minima:
 $(\kappa_V, \kappa_f) = (1.05, -2.40)$ and
 $(\kappa_V, \kappa_f) = (1.05, 2.30)$
- The integral of the posterior density in the $(+,+)$ quadrant is 26% of the total, while the remaining 74% of the integral of the posterior density is contained within the $(+,-)$ quadrant



Summary on couplings


- Couplings to fermions: $\kappa_f = -2.64^{+1.59}_{-1.30}$
- Couplings to bosons:
 $\kappa_W = -1.27^{+0.46}_{-0.29}$; *second interval* $1.04 < \kappa_W < 1.51$
 $\kappa_Z = \pm 1.05^{+0.45}_{-0.55}$
 - if varied together: $(\kappa_W, \kappa_Z) = (1.25, \pm 0.90)$
- For custodial symmetry: $|\theta_{WZ}| = 0.68^{+0.21}_{-0.41} \rightarrow \lambda_{WZ} = 1.24^{+2.34}_{-0.42}$
- If custodial symmetry is preserved:
 $(\kappa_V, \kappa_f) = (1.05, -2.40)$ and $(\kappa_V, \kappa_f) = (1.05, 2.30)$

Summary

- Tevatron has ended its 25 years' run on September 30th 2011
 - It ran more than 9 years at $\sqrt{s} = 1.96$ TeV
 - It delivered almost 12 fb^{-1} during that period
 - We are grateful for all these data

- D0 finalized publications of all search channels

- Broad data excess compatible with a Higgs boson with $m_H = 125 \text{ GeV}$ is observed in the two main channels, $VH \rightarrow Vbb$ and $H \rightarrow WW$, as well as in the full combination





- All results just submitted for publication:



http://www-d0.fnal.gov/d0_publications/d0_pubs_list_bytopic.html#higgs




Summary

- Both CDF and D0 experiments performed very well
 - Improvements over the years led to the 95% C.L. exclusion sensitivity $< \sim 1.0 \times \text{SM}$ for $m_H < 185 \text{ GeV}$ when combining two experiments
- 3.1 s.d. excess @125 GeV observed in data when combining from both experiments, consistent with LHC observation
- Signal strengths in all analyzed decay channels are consistent with SM Higgs expectation
- Results on Higgs couplings are also consistent with the SM predictions
- It is unlikely that $H \rightarrow b\bar{b}$ is established before 2015, except if the results from all experiments are combined



Backup



Signal strength for various decays



Tevatron $H \rightarrow b\bar{b}$ Results

PRL 109,071804(2012)

- Last Summer:
 - $\sigma_{VH} = 0.23 \pm 0.09$ pb (SM: 0.12 ± 0.01 pb) @125 GeV
- Now:
 - $\sigma_{VH} = 0.19 \pm 0.09$ pb, consistent with the summer results
 - The shift in this result is due to the updated $ZH \rightarrow \nu\nu b\bar{b}$ analysis from CDF and corresponds to a change in the central value of 0.6 times the total uncertainty, consistent with the difference expected given the observed changes in the CDF $ZH \rightarrow \nu\nu b\bar{b}$





CDF results presented by T. Junk on January 18th:

http://theory.fnal.gov/jetp/talks/wc_trj_cdfhiggs_18jan_pub.pdf

Interpretation in non SM





• α

• α



H \rightarrow $\tau\tau$

• a



The Higgs Mechanism

- Essential ingredient of the **Standard Model**
 - Complex scalar field with potential
- Used to **break the el. weak symmetry**...


$$M_W = \frac{1}{2} v g \quad M_Z = \frac{1}{2} v g / \cos \theta_W = M_W / \cos \theta_W$$

- ... and to **generate fermion masses**:

$$m_f = g_f v / \sqrt{2} \Rightarrow g_f = m_f \sqrt{2} / v$$

- Unitarity requires a scalar Higgs boson or similar

- **cross section for WW scattering diverges like s/M_W^2**
- **scalar Higgs boson cancels divergences**





Systematics

- Luminosity: 6.1%
- b-tagging rate: 1-10%
- JES and JER $\sim 7\%$
- Lepton id and similar: 1-9%
- Simulated backgrounds
cross sections 4-30%
- MJ background 10-30%

CDF and DØ experiments in Run II

- Both detectors are upgraded in Run II
- New silicon micro-vertex trackers
- New tracking systems
- Upgraded muon chambers





Angular coverage	$ \eta $
Muon ID	~ 2
Tracking	~ 2.5
EM / Jet ID	~ 4

An example of limits settings

- Compare Poisson likelihood of B hypothesis to S+B hypothesis, and calculate their negative log likelihood ratio (LLR):

$L(B)$	$L(S+B)$	LLR
$\prod_i \frac{b_i^{d_i} \exp(-b_i)}{d_i!}$	$\prod_i \frac{(s_i + b_i)^{d_i} \exp(-(s_i + b_i))}{d_i!}$	$2 \cdot \sum_i s_i - d_i \cdot \log(1 + s_i/b_i)$

- where d_i events observed in bin i with S and B expectations s_i and b_i .
- or b_i in each bin (B) or $s_i + b_i$ in each bin (S+B)
- Repeat many times to obtain LLR distribution: median is Expected LLR



S+B p-values

